

Chemical Constituents of *Verbascum L.* Species

İ. İrem TATLI*^o, Zeliha Ş. AKDEMİR*

Chemical Constituents of *Verbascum L.* Species

Summary

Verbascum species (Scrophulariaceae), which are widespread plants in Anatolia, are used as expectorant and mucolytic in folk medicine. In this paper, the chemical constituents of saponins, iridoid and phenylethanoid glycosides, monoterpene glucoside, neolignan glucosides, flavonoids, steroids, spermine alkaloids and other compounds of these species are reviewed.

Key words : *Verbascum L. species*, Scrophulariaceae, mullein.

Received : 07.06.2004

Revised : 25.10.2004

Accepted : 04.11.2004

Verbascum Türlerinin Kimyasal Bileşimi

Özet

Anadolu'da çok yaygın olarak yetişen *Verbascum türleri* (Scrophulariaceae) halk arasında balgam söktürücü ve göğüs yumuşatıcı olarak kullanılmaktadır. Bu derlemede, türlerin kimyasal bileşiminde yer alan saponinler, iridoit ve feniletanoit glikozitleri, monotermen glukoziti, neolignan glukozitleri, flavonoitler, steroidler, spermin alkaloitleri ve diğer bileşikler hakkında bilgi verilmiştir.

Anahtar Kelimeler: Sığırkuyruğu, *Verbascum L. türleri*, Scrophulariaceae, mullein.

INTRODUCTION

Verbascum L. is the largest genus of the family Scrophulariaceae, with about 2500 species worldwide¹. This genus, commonly known as "Sığırkuyruğu", is represented by 228 species, of which 185 are endemic in the flora of Turkey². Various preparations of some species of this genus have been used as expectorant and mucolytic, as well as sudorific, sedative, diuretic and constipate in traditional Turkish medicine³. *Verbascum species* are also used externally for desiccating wounds, anal fistula and pruritic conditions in urogenital organs⁴. Additionally, in Europe, Asia and Northern America, several *Verbascum species* have been reported as antiseptic,

astringent, demulcent, emollient, expectorant, sedative, narcotic, diuretic and antimalarial and as a treatment for tumors, inflammations, migraine, asthma and spasmodic coughs¹.

We present here an exhaustive review of the literature, available as of the end of 2004, on the metabolites from the *Verbascum species* throughout the world.

The compounds in *Verbascum spec.* are classified as eight main groups, such as saponins, iridoid and phenylethanoid glycosides, monoterpene glucoside, neolignan glucosides, flavonoids, steroids and spermine alkaloids. Phenolic acids, fatty acids and some


* Hacettepe University, Faculty of Pharmacy, Department of Pharmacognosy, 06100, Ankara TURKEY

^o Corresponding author e-mail: itatli@hacettepe.edu.tr

other compounds are included under the title, "Other Compounds".


The major skeleton of each main group is given in the relevant section. The structures are represented in the tables from the basic to the complex skeleton, together with distribution in plants in alphabetical order (*Recorded species in the flora of Turkey).

I. Saponins:




This is one of the major groups in the genus *Verbascum*. The phytochemical studies on these species have revealed the presence of oleanane type triterpene saponins, usually monodesmosidic, and their sugar moieties are attached at C-3 (OH). The saponins in *Verbascum* species are given in Table 1.

II. Iridoid Glycosides:




Iridoids, which represent a large group of cyclopentano[c]pyran monoterpenoids, are found as natural constituents in a large number of *Verbascum* species. A bicyclic H-5/H-9 β,β -cis-fused cyclopentapyran ring system is the most common structural feature of these substances. They also contain a double bond between C-3 and C-4 including non-substitution at C-4. *Verbascum* species contain especially aucubin and catalpol type iridoids and their acylated derivatives are modified with aliphatic and aromatic acids (e.g. acetic acid, *trans*-cinnamic acid, *trans-p*-coumaric acid, ferulic acid); these glycosides are also varied with the esterification positions. The iridoid glycosides in *Verbascum* species are shown in Table 2.

III. Phenylethanoid Glycosides:




Phenylethanoid glycosides are natural polyphenolic compounds widely seen in many Dicotyledone families, and are also rich in the *Verbascum* genus. Structurally, they are characterized as glycosides of phenylethanol, esterified at R₁ position by cinnamic, caffeic and ferulic acid derivatives. Attachment of the increasing number of sugar units (to date, mono, di, tri), the type of the sugar (glucose, rhamnose, arabinose, apiose and xylose), different sequence (R₂-R₄) and substitutions cause great variations in their structures. The phenylethanoid glycosides in *Verbascum* species are given in Table 3.

IV. Monoterpene Glucosides:



In a continuation of the phytochemical studies on *Verbascum* species of Turkish origin, a monoterpene glucoside, 1- β -glucopyranosyl-8-hydroxy-3,7-dimethyl-2(*E*),6(*E*)-octadienoate, was isolated from the aerial parts of *V. pterocalycinum* var. *mutense*⁹.

V. Neolignan Glucosides:



Lignans and neolignans belong to an important group of natural products, consisting of two phenylpropane monomers linked through carbon-carbon or carbon-oxygen bonds. The neolignan glucosides

Table 1. Saponins of *Verbascum* L. species

Saponins	Species	References
Saponin II	<i>V. nigrum</i>	5
Verbascosaponin (= Saponin I)	<i>V. nigrum</i>	5
	<i>V. phlomoides</i> *	6
Songarosaponin B	<i>V. songaricum</i> *	7
Ilwensisaponin C	<i>V. nigrum</i>	8
	<i>V. pterocalycinum</i> var. <i>mutense</i> *	9
Thapsuine B	<i>V. lychnitis</i>	10
	<i>V. thapsus</i> *	11
Hydroxythapsuine B	<i>V. thapsus</i> *	11
Triterpene B	<i>V. phlomoides</i> *	6
	<i>V. thapsus</i> *	12
Triterpene A	<i>V. phlomoides</i> *	6
	<i>V. thapsus</i> *	12
Saikogenin A	<i>V. thapsus</i> *	12
Songarosaponin A	<i>V. songaricum</i> *	7
Songarosaponin E	<i>V. songaricum</i> *	13
Songarosaponin F	<i>V. songaricum</i> *	13
3- <i>O</i> - β -D-fucopyranosylsaikogenin F	<i>V. roripifolium</i>	14
Mulleinsaponin I	<i>V. sinaiticum</i>	14
Saikosaponin A	<i>V. roripifolium</i>	14
	<i>V. sinaiticum</i>	
Mulleinsaponin II	<i>V. sinaiticum</i>	14
Mulleinsaponin III	<i>V. roripifolium</i>	14
	<i>V. sinaiticum</i>	
Desrhamnosyl verbascosaponin	<i>V. phlomoides</i> *	15
	<i>V. thapsiforme</i> *	14
Songarosaponin C	<i>V. songaricum</i> *	7
	<i>V. thapsiforme</i> *	14
Songarosaponin D	<i>V. songaricum</i> *	16

	<i>V. thapsiforme</i> *	14
Buddlejasaponin IV	<i>V. thapsiforme</i> *	14
İlwensisaponin A (= Mimengoside A)	<i>V. pterocalycinum</i> var. <i>mutense</i> *	9
	<i>V. roripifolium</i>	14
	<i>V. sinaiticum</i>	
	<i>V. thapsiforme</i> *	8
	<i>V. nigrum</i>	14
Buddejasaponin I (= Verbascosaponin B)	<i>V. fruticosum</i>	14
	<i>V. roripifolium</i>	
	<i>V. sinaiticum</i>	
	<i>V. songaricum</i> *	13
	<i>V. thapsiforme</i> *	14
Mulleinsaponin IV	<i>V. fruticosum</i>	14
Mulleinsaponin V	<i>V. fruticosum</i>	14
Mulleinsaponin VI	<i>V. sinaiticum</i>	14
Mulleinsaponin VII	<i>V. sinaiticum</i>	14
3-O-[[β -D-glucosyl-(1 \rightarrow 2)- β -D-fucopyranosyl-(1 \rightarrow 4)]- α -L-rhamnopyranosyl]-(1 \rightarrow 2)- β -D-glucopyranosyl]-13 β ,28-epoxy-olean-11-ene-3 β ,23-diol	<i>V. lychnitis</i>	
3-O-[[β -D-glucosyl-(1 \rightarrow 2)- β -D-fucopyranosyl-(1 \rightarrow 4)]- α -L-rhamnopyranosyl]-(1 \rightarrow 2)- β -D-glucopyranosyl]-16 β -(β -D-glucopyranosyloxy)-13 β ,28-epoxy-olean-11-ene-3 β , 23-diol	<i>V. lychnitis</i>	17
Thapsuine A	<i>V. lychnitis</i>	10
	<i>V. thapsus</i> *	11
Hydroxythapsuine A	<i>V. thapsus</i> *	11
Niga-ichigoside F1	<i>V. wiedemannianum</i> *	18
Rosmatin	<i>V. wiedemannianum</i> *	18
Ursolic acid	<i>V. lychnitis</i>	10
Ancrytosaponin A	<i>V. ancyritanum</i> *	19

* Recorded species in the flora of Turkey

Table 2. Iridoid glycosides of *Verbascum* L. species

Iridoid glycosides	Species	References
Aucubin	<i>V. cheirantifolium</i> *	20
	<i>V. georgicum</i> *	21
	<i>V. lasianthum</i> *	22
	<i>V. laxum</i>	23
	<i>V. lychnitis</i>	10
	<i>V. nigrum</i>	24
	<i>V. phlomoides</i> *	25
	<i>V. saccatum</i> *	26
	<i>V. sinuatum</i> *	27
	<i>V. spinosum</i>	28
	<i>V. thapsiforme</i> *	24
	<i>V. thapsus</i> *	24
	<i>V. undulatum</i>	29
	<i>V. virgatum</i>	30
<i>V. wiedemannianum</i> *	18	
6- <i>O</i> - β -D-glucopyranosyl aucubin	<i>V. sinuatum</i> *	31
Sinuatol	<i>V. laxum</i>	23
	<i>V. nigrum</i>	32
	<i>V. sinuatum</i> *	33
	<i>V. undulatum</i>	34
6- <i>O</i> - β -D-xylopyranosyl aucubin	<i>V. sinuatum</i> *	35
	<i>V. thapsiforme</i> *	36
	<i>V. thapsus</i> *	27
6- <i>O</i> - α -L-sinuatosyl aucubin	<i>V. sinuatum</i> *	37
Sinuatocide	<i>V. sinuatum</i> *	37
Nigroside II	<i>V. nigrum</i>	32
	<i>V. undulatum</i>	34
Nigroside I	<i>V. nigrum</i>	32
	<i>V. undulatum</i>	29
Nigroside III	<i>V. nigrum</i>	38
	<i>V. undulatum</i>	34
6- <i>O</i> -(3''- <i>O</i> - <i>p</i> -coumaroyl)- α -L-rhamnopyranosyl aucubin	<i>V. laxum</i>	23
	<i>V. nigrum</i>	38
	<i>V. undulatum</i>	29
Unduloside III	<i>V. lasianthum</i> *	22
	<i>V. undulatum</i>	39
Unduloside II	<i>V. undulatum</i>	39
6- <i>O</i> -(2''- <i>O</i> - <i>p</i> -methoxy- <i>trans</i> -cinnamoyl, 3''- <i>O</i> -acetyl)- α -L-rhamnopyranosylaucubin	<i>V. laxum</i>	23
Phlomoidoside	<i>V. phlomoides</i> *	15
Unduloside	<i>V. undulatum</i>	34
6- <i>O</i> -(3''- <i>O</i> - <i>trans</i> -feruloyl)- α -L-rhamnopyranosylaucubin	<i>V. undulatum</i>	29
Nigroside V	<i>V. nigrum</i>	38
Nigroside IV	<i>V. nigrum</i>	38
6- <i>O</i> -(<i>p</i> -methoxy-cinnamoyl)-aucubin	<i>V. cheirantifolium</i> *	20
6- <i>O</i> -(<i>p</i> -coumaroyl) aucubin	<i>V. cheirantifolium</i> *	20
Aucuboside	<i>V. chaixii</i>	40
	<i>V. lychnitis</i>	41

	<i>V. sinuatum*</i>	27
	<i>V. phlomooides*</i>	42
	<i>V. thapsiforme*</i>	40
Catalpol	<i>V. chaixii</i>	40
	<i>V. cheirantifolium*</i>	20
	<i>V. cilicicum*</i>	43
	<i>V. georgicum*</i>	21
	<i>V. laxum</i>	23
	<i>V. lychnitis</i>	41
	<i>V. nigrum</i>	24
	<i>V. phlomooides*</i>	25
	<i>V. sinuatum*</i>	44
	<i>V. spinosum</i>	28
	<i>V. thapsiforme*</i>	42
	<i>V. thapsus*</i>	24
	<i>V. wiedemannianum*</i>	18
Isocatalpol	<i>V. lychnitis</i>	41
	<i>V. nigrum</i>	24
	<i>V. thapsiforme*</i>	
	<i>V. thapsus*</i>	
Methylcatalpol	<i>V. lychnitis</i>	41
	<i>V. nigrum</i>	24
	<i>V. thapsiforme*</i>	
	<i>V. thapsus*</i>	
6- <i>O</i> - α -L-rhamnopyranosylcatalpol	<i>V. georgicum*</i>	21
	<i>V. lasianthum*</i>	22
	<i>V. saccatum*</i>	26
	<i>V. thapsus*</i>	45
6- <i>O</i> - β -D-xylopyranosylcatalpol	<i>V. thapsiforme*</i>	35
Verbaspinoside	<i>V. cilicicum*</i>	43
	<i>V. spinosum</i>	28
6- <i>O</i> -(3''- <i>O</i> - <i>trans</i> -cinnamoyl)- α -L-rhamnopyranosylcatalpol	<i>V. cilicicum*</i>	43
6- <i>O</i> -(4''- <i>O</i> - <i>trans</i> -cinnamoyl)- α -L-rhamnopyranosylcatalpol	<i>V. cilicicum*</i>	43
Saccatoside	<i>V. cilicicum*</i>	43
	<i>V. saccatum*</i>	26
	<i>V. thapsus*</i>	45
	<i>V. virgatum</i>	30
6- <i>O</i> -(3''- <i>O</i> - <i>p</i> -coumaroyl)- α -L-rhamnopyranosylcatalpol	<i>V. sinuatum*</i>	44
	<i>V. thapsus*</i>	45
	<i>V. virgatum</i>	30
6- <i>O</i> -(4''- <i>O</i> - <i>p</i> -coumaroyl)- α -L-rhamnopyranosylcatalpol	<i>V. thapsus*</i>	45
6- <i>O</i> -(2''- <i>O</i> -(<i>p</i> -methoxy- <i>trans</i> -cinnamoyl))- α -L-rhamnopyranosylcatalpol	<i>V. thapsus*</i>	45
6- <i>O</i> -(3''- <i>O</i> -(<i>p</i> -methoxy- <i>trans</i> -cinnamoyl))- α -L-rhamnopyranosylcatalpol	<i>V. thapsus*</i>	45
Verbascoside A	<i>V. georgicum*</i>	21
	<i>V. lasianthum*</i>	22
	<i>V. thapsus*</i>	45
6- <i>O</i> -[2''- <i>O</i> -(3,4-dihydroxy- <i>trans</i> -cinnamoyl)]- α -L-rhamnopyranosylcatalpol	<i>V. thapsus*</i>	45
6- <i>O</i> -[4''- <i>O</i> -(3,4-dihydroxy- <i>trans</i> -cinnamoyl)]- α -L-rhamnopyranosylcatalpol	<i>V. thapsus*</i>	45
6- <i>O</i> -[3''- <i>O</i> -(3,4-dimethoxy- <i>trans</i> -cinnamoyl)]- α -L-	<i>V. thapsus*</i>	45

rhamnopyranosylcatalpol		
6-O-(2''-O-feruloyl)- α -L-rhamnopyranosylcatalpol	<i>V. thapsus</i> *	45
6-O-(4''-O-feruloyl)- α -L-rhamnopyranosylcatalpol	<i>V. thapsus</i> *	45
6-O-(2''-O-isoferuloyl)- α -L-rhamnopyranosylcatalpol	<i>V. thapsus</i> *	45
6-O-(3''-O-isoferuloyl)- α -L-rhamnopyranosylcatalpol	<i>V. thapsus</i> *	45
6-O-(4''-O-isoferuloyl)- α -L-rhamnopyranosylcatalpol	<i>V. thapsus</i> *	45
Pulverulentoside I	<i>V. lasianthum</i> *	22
	<i>V. laxum</i>	23
	<i>V. pulverulentum</i>	46
	<i>V. sinuatum</i> *	47
	<i>V. thapsus</i> *	45
6-O-(2''-O- <i>p</i> -methoxy- <i>trans</i> -cinnamoyl-4''-O-asetyl)- α -L-rhamnopyranosylcatalpol	<i>V. thapsus</i> *	45
Buddlejoside A ₅	<i>V. lasianthum</i> *	22
Pulverulentoside II	<i>V. pulverulentum</i>	49
Catalposide	<i>V. chaixii</i>	40
	<i>V. lychnitis</i>	41
6-O-(<i>p</i> -coumaroyl) catalpol (= Specioside)	<i>V. lychnitis</i>	48
	<i>V. phlomoides</i> *	15
Picoside IV	<i>V. pterocalycinum</i> var. <i>mutense</i> *	9
Angeloside	<i>V. wiedemannianum</i> *	18
Ajugol	<i>V. pterocalycinum</i> var. <i>mutense</i> *	9
	<i>V. spinosum</i>	28
	<i>V. thapsus</i> *	45
	<i>V. undulatum</i>	29
	<i>V. virgatum</i>	30
	<i>V. wiedemannianum</i> *	18
6-O-benzoyl ajugol	<i>V. thapsus</i> *	45
6-O-syringoyl ajugol	<i>V. thapsus</i> *	45
6-O-vanilloyl ajugol	<i>V. lasianthum</i> *	49
	<i>V. thapsus</i> *	45
Harpagide	<i>V. phlomoides</i> *	42
	<i>V. sinuatum</i> *	27
	<i>V. thapsiforme</i> *	42
	<i>V. thapsus</i> *	31
	<i>V. undulatum</i>	50
8-O-acetyl harpagide	<i>V. lasianthum</i> *	49
	<i>V. phlomoides</i> *	42
	<i>V. thapsiforme</i> *	
Verbascoside B	<i>V. georgicum</i> *	51
Harpagoside	<i>V. lasianthum</i> *	49
	<i>V. laxum</i>	23
	<i>V. nigrum</i>	32
	<i>V. phlomoides</i> *	42
	<i>V. pulverulentum</i>	46
	<i>V. sinuatum</i> *	47
	<i>V. thapsiforme</i> *	42
	<i>V. thapsus</i> *	45
	<i>V. undulatum</i>	50
Geniposidic acid	<i>V. olympicum</i> *	52
Lychnitoside	<i>V. lychnitis</i>	10
Glutinoside	<i>V. wiedemannianum</i> *	18
Rehmaglutin D	<i>V. wiedemannianum</i> *	18

* Recorded species in the flora of Turkey

Table 3. Phenylethanoid glycosides of *Verbascum* L. species

Phenylethanoid glycosides	Species	References
Verbascoside (=Acteoside)	<i>V. cilicicum</i> *	43
	<i>V. georgicum</i> *	53
	<i>V. lasianthum</i> *	49
	<i>V. lychnitis</i>	54
	<i>V. nigrum</i>	54
	<i>V. phlomoides</i> *	54
	<i>V. pterocalycinum</i> var. <i>mutense</i> *	9
	<i>V. salviifolium</i> *	55
	<i>V. sinaiticum</i>	56
	<i>V. sinuatum</i> *	57
	<i>V. spinosum</i>	28
	<i>V. thapsiforme</i> *	54
	<i>V. undulatum</i>	34
<i>V. wiedemannianum</i> *	58	
β -Hydroxyacteoside	<i>V. salviifolium</i> *	55
2'-O-Acetylacteoside	<i>V. sinaiticum</i>	59
Martynoside	<i>V. salviifolium</i> *	55
	<i>V. undulatum</i>	29
	<i>V. wiedemannianum</i> *	58
6'-O-Acetyl-martynoside (= Wiedemannioside A)	<i>V. undulatum</i>	29
	<i>V. wiedemannianum</i> *	58
Echinacoside	<i>V. wiedemannianum</i> *	58
Poliumoside	<i>V. blattaria</i> *	60
	<i>V. boerhavii</i>	
	<i>V. chaixii</i>	
	<i>V. lasianthum</i> *	49
	<i>V. phlomoides</i> *	60
	<i>V. sinuatum</i> *	
	<i>V. thapsus</i> *	
Forsythoside B	<i>V. lychnitis</i>	61
	<i>V. nigrum</i>	61
	<i>V. phlomoides</i> *	15
	<i>V. salviifolium</i> *	55
	<i>V. thapsiforme</i> *	61
	<i>V. thapsus</i> *	62
Arenarioside	<i>V. sinaiticum</i>	59
	<i>V. thapsus</i> *	62
	<i>V. undulatum</i>	29
Alyssonoside	<i>V. thapsus</i> *	62
1'-O- β -D-(3,4-dihydroxy-phenyl)-ethyl- α -L-rhamnopyranosyl-(1 \rightarrow 3')- β -D-xylopyranosyl-(1 \rightarrow 6')-4'-O-feruloyl-glucopyranoside	<i>V. sinaiticum</i>	59
	<i>V. thapsus</i> *	62
Angoroside A	<i>V. salviifolium</i> *	55
	<i>V. spinosum</i>	28
Angoroside C	<i>V. spinosum</i>	28
Leucosceptoside B	<i>V. thapsus</i> *	62
	<i>V. wiedemannianum</i> *	58
1'-O- β -D-(3-hydroxy-4-methoxy-phenyl)-ethyl- α -L-rhamnopyranosyl-(1 \rightarrow 3')- β -D-xylopyranosyl-(1 \rightarrow 6')-4'-O-feruloyl-glucopyranoside	<i>V. thapsus</i> *	62


Cistanoside B	<i>V. sinaiticum</i>	59
	<i>V. thapsus</i> *	62
Wiedemannioside B	<i>V. wiedemannianum</i> *	58
Wiedemannioside C	<i>V. wiedemannianum</i> *	58
Wiedemannioside D	<i>V. wiedemannianum</i> *	58
Wiedemannioside E	<i>V. wiedemannianum</i> *	58
1'- <i>O</i> - β -D-(3,4-dihydroxy-phenyl)-ethyl- α -L-rhamnopyranosyl- (1 \rightarrow 3')-3'''-hydroxy-4'''- <i>O</i> - β -D-glucopyranosyl-cinnamoyl-(1 \rightarrow 6')-glucopyranoside	<i>V. thapsus</i> *	62

* Recorded species in the flora of Turkey

from *Verbascum* species have dehydrodiconiferyl alcohol skeleton.

So far, two *Verbascum* species, *Verbascum thapsus*⁶² and *V. salviifolium*⁵⁵, have been reported to contain neolignan glucosides, such as dehydrodiconiferyl alcohol 9-*O*- β -D-glucopyranoside (7*R*, 8*S*), dehydrodiconiferyl alcohol 9-*O*- β -D-glucopyranoside (7*S*, 8*R*), 5-*O*-methyldehydrodiconiferyl alcohol 9-*O*- β -D-glucopyranoside (7*R*, 8*S*), 5-*O*-methyldehydrodiconiferyl alcohol 9-*O*- β -D-glucopyranoside (7*S*, 8*R*), 4-*O*- β -D-glucopyranosyl-dehydrodiconiferyl alcohol 9-*O*- β -D-glucopyranoside (7*R*, 8*S*) and dehydrodiconiferyl alcohol 9'-*O*- β -D-glucopyranoside.


VI. Flavonoids:



The flavonoid glycosides are widely distributed in the plant kingdom. They have benzo- γ -pyrone skeleton in their structure. To date, many flavonoids have been reported from the *Verbascum* species, in which there was a large variety of flavone and flavonol aglycones, such as apigenin, luteolin, quercetin and kaempferol. Glycosylation was usually at C-7 position of these aglycones. On the other hand, a C-

glucoside was found in *V. cherianthifolium*. Isoflavone and flavonolignans from *V. sinaiticum* have also been observed. The flavonoids from *Verbascum* species are seen in Table 4.

VII. Steroids:



Khuroo and his colleagues determined steroid derivatives in *Verbascum* species³⁶. Structurally, these compounds usually contain cholestanon skeleton. 24 α -methyl-5 α -cholestan-3-on, 24 α -ethyl-5 α -cholestan-3-on, 24 α -ethyl-5 β -cholestan-3-on, 24 α -ethyl-5 α -cholestan-7-en-3-on, 24 α -ethyl-5 α -cholestan-22-en-3-on, 24 α -ethyl-5 β -cholestan-22-en-3-on, 24 α -ethyl-5 α -cholestan- $\Delta^{7,22}$ -dien-3-on and β -sitosterol were isolated from *V. thapsus*³⁶. β -sitosterol was also obtained from the aerial parts of *V. lasianthum*⁷⁵, *V. phlomoides*²⁵, *V. pycnostachyum*³⁷⁶ and *V. thapsiforme*⁷⁷. Moreover, sitosterol-3-*O*- β -D-glucopyranoside¹⁸ from *V. wiedemannianum*, stigmasterol from *V. pycnostachyum*⁷⁶ and *V. phlomoides*²⁵, phytosterol from *V. phlomoides*⁷⁷ and *V. thapsiforme*⁷⁷, and phytosterol glucoside from *V. phlomoides*⁷⁸ were reported.


Table 4. Flavonoids of *Verbascum* L. species

Flavonoids	Species	References
Apigenin	<i>V. cheiranthifolium</i> *	63
	<i>V. eremobium</i>	64
	<i>V. fruticosum</i>	64
	<i>V. lychnitis</i>	41
	<i>V. phlomoides</i> *	65
	<i>V. schimperianum</i>	64
	<i>V. songaricum</i> *	66
	<i>V. thapsiforme</i> *	67
Apigenin-7-glucoside	<i>V. eremobium</i>	64
	<i>V. fruticosum</i>	64
	<i>V. letourneuii</i>	64
	<i>V. phlomoides</i> *	65
	<i>V. salviifolium</i> *	55
	<i>V. schimperianum</i>	64
	<i>V. siniaticum</i>	64
	<i>V. thapsiforme</i> *	67
Apigenin-7-glucuronide	<i>V. lychnitis</i>	68
	<i>V. nigrum</i>	
Apigenin-4'-rhamnoside	<i>V. thapsus</i> subsp. <i>thapsus</i> *	69
Luteolin	<i>V. cheiranthifolium</i> *	63
	<i>V. lychnitis</i>	41
	<i>V. fruticosum</i>	64
	<i>V. phlomoides</i> *	65
	<i>V. siniaticum</i>	70
	<i>V. songaricum</i> *	66
	<i>V. thapsiforme</i> *	67
	<i>V. thapsus</i> *	71
	<i>V. wiedemannianum</i> *	18
Luteolin-5-glucoside	<i>V. lychnitis</i>	41
Luteolin-7- glucoside	<i>V. eremobium</i>	64
	<i>V. fruticosum</i>	64
	<i>V. letourneuii</i>	64
	<i>V. phlomoides</i> *	65
	<i>V. salviifolium</i> *	55
	<i>V. scardicolum</i>	72
	<i>V. schimperianum</i>	64
	<i>V. siniaticum</i>	64
	<i>V. songaricum</i> *	66
	<i>V. thapsiforme</i> *	67
Luteolin-3'- glucoside	<i>V. salviifolium</i> *	55
Luteolin-7-glucuronide	<i>V. lychnitis</i>	68
	<i>V. nigrum</i>	
6-hydroxy-luteolin-7- glucoside	<i>V. thapsus</i> subsp. <i>thapsus</i> *	69
7-methoxy-luteolin	<i>V. lychnitis</i>	41
Verbacoside	<i>V. thapsus</i> *	71
Acacetin	<i>V. lychnitis</i>	41
Acacetin-7- glucoside	<i>V. fruticosum</i>	64
	<i>V. schimperianum</i>	
	<i>V. siniaticum</i>	
Acacetin-7-galactoside	<i>V. fruticosum</i>	64
	<i>V. schimperianum</i>	
	<i>V. siniaticum</i>	
Diosmin	<i>V. phlomoides</i> *	73

Hesperidin	<i>V. phlomoides</i> *	73
Diosmetin-7-glucuronide	<i>V. nigrum</i>	68
Chrysoeriol	<i>V. phlomoides</i> *	65
	<i>V. siniaticum</i>	70
Chrysoeriol-7-glucoside	<i>V. eremobium</i>	64
	<i>V. fruticosum</i>	
	<i>V. salviifolium</i> *	55
	<i>V. schimperianum</i>	64
	<i>V. siniaticum</i>	
Chrysoeriol-7-diglucoside	<i>V. eremobium</i>	64
	<i>V. fruticosum</i>	
	<i>V. letourneuii</i>	
	<i>V. siniaticum</i>	
7, 4'-dihydroxy flavon-4-rhamnoside	<i>V. thapsus</i> *	69
Vitexin	<i>V. cheiranthifolium</i> *	63
Swertisin	<i>V. cheiranthifolium</i> *	63
Kaemferol	<i>V. phlomoides</i> *	65
Quercetin	<i>V. lychnitis</i>	68
	<i>V. phlomoides</i> *	65
	<i>V. songaricum</i> *	66
Quercetin-7-glucoside	<i>V. thapsiforme</i> *	67
Quercetin-7-glucuronide	<i>V. lychnitis</i>	68
	<i>V. nigrum</i>	
Quercetin-3,7-diglucoside	<i>V. thapsiforme</i> *	67
Rutin	<i>V. phlomoides</i> *	65
İsorhamnetin	<i>V. thapsus subsp. thapsus</i> *	69
Tamarixetin-7-glucoside	<i>V. phlomoides</i> *	73
Tamarixetin-7-rutinoside	<i>V. phlomoides</i> *	73
5-hydroxy-6,7- dimethoxyflavone-3-ol	<i>V. thapsus</i> *	74
Patuletin	<i>V. lychnitis</i>	41
Orobol	<i>V. siniaticum</i>	56
Orobol-7-glucoside	<i>V. siniaticum</i>	56
8-methylorobol-7-glucoside	<i>V. siniaticum</i>	56
3', 4'-dimethoxy-orobol-7- rhamnoside	<i>V. siniaticum</i>	56
Eriodictiol	<i>V. phlomoides</i> *	65
Sinaiticin	<i>V. sinaiticum</i>	70
Hydrocarpin	<i>V. sinaiticum</i>	70

* Recorded species in the flora of Turkey

VIII. Spermine Alkaloids:



The isolation and structural elucidation of the novel 17-membered lactam alkaloids, verbacine and verballocine, were reported from *Verbascum pseudonobile* Stoj. et Stef.⁷⁹, together with their corresponding *N,N'*-methylene-bridge derivatives, namely (+)-(*S*)-verbamethine, (+)-(*S*)-isoverbamethine, (-)-verbasitrine, (-)-isoverbasitrine, (+)-verbametrine and (+)-isoverbametrine⁸⁰. Verbaskine was also isolated from this plant, for which evidence of the artificial origin was previously presented⁷⁹.

Furthermore, (*S*)-verbasikrin, (*S*)-isoverbasikrin, (*S*)-verbamekrin and (*S*)-isoverbamekrin were obtained from *V. pseudonobile* and structurally characterized⁸¹. In the same plant material, the presence of the spermine alkaloids, protoverbine (8*S*)-8-phenyl-1,5,9,13-tetraazacycloheptadecane-6-on, protometin (2*S*)-2-phenyl-1,5,9,14-tetraazabicyclo [12.3.1]-octadecane-4-on⁸², verbamedine, isoverbamedine (*Z*) and *N*(13)-formimino-verbacine were reported⁸³. Verbacine and verbascenine were also found in *V. nobile*⁸⁴. Verbascenine and verballoscenine were yielded from *V. phoeniceum* L. Verbascenine was also elucidated from the total alkaloid extracts of *V. nigrum* L.⁸⁵.

IX. Other Compounds:

The presence of anabasine and plantagonine alkaloids from the leaves of *V. songaricum* Schrenk was determined⁸⁶. 3 α -hydroxydrimmanyl-8-methanoate, a sesquiterpene, was reported from *V. thapsus*³⁶. Magiatis and his colleagues isolated verbalactone, a macrocyclic dimer lactone, from the roots of *V. undulatum*⁵⁰.

V. phlomoides contained phenolic acids as dihydroxy cinnamic acid, *para*-coumaric acid, ferulic acid,

para-hydroxy benzoic acid and vanillic acid^{25,87}. However the contents of *para*-coumaric acid, ferulic acid, *para*-hydroxy benzoic acid and vanillic acid in *V. thapsiforme* were two times more than their amounts in *V. phlomoides*⁸⁷.

The benzene extract of seed oil of *V. thapsus* contained palmitic, stearic, oleic, linoleic, linolenic, arachidic and behenic acids in the nonsaponifiable fraction⁸⁸. *V. phlomoides* and *V. thapsiforme* were also rich in palmitic and linoleic acids⁸⁷.

The water-soluble polysaccharides of mullein flowers (*Verbascum*) were isolated and characterized. The neutral fraction consisted of a xyloglucan and an arabinogalactan⁸⁹.

RESULTS and DISCUSSION

As a result of exhaustive investigations, it is noticed that only 40 species have been studied, 16 of which are recorded in the flora of Turkey.

Verbascum species have been used as expectorant, mucolytic and diuretic as well as in the treatment of hemorrhoids.

The isolated compounds from *Verbascum* species were evaluated under eight main groups as: saponins, iridoids, phenylethanoids, monoterpene and neolignan glycosides, flavonoids, steroids and spermine alkaloids. Most of the compounds were isolated from the aerial parts of the plants.

It is considered that iridoid glycosides, flavonoids, phenylethanoid glycosides and saponins, respectively, are obtained in the plants more than the others and that they may be responsible for the biological activities mentioned above.

V. thapsus, which has officinal preparations, is also rich in the mentioned compounds. In our ongoing project, developing officinal preparations from some endemic *Verbascum* species will be useful. Moreover, the chemical characterization of *Verbascum* species growing in Turkey will be determined.

REFERENCES

- Grieve M. A Modern Herbal, Barnes and Noble Books, New York, 564-6, 1995.
- Huber-Morath A. *Verbascum* L., Davis PH (ed.), Flora of Turkey and the East Aegean Islands, University Press, Edinburgh, Vol. 6, 461-603, 1978.
- Baytop T. Therapy with Medicinal Plants in Turkey (Past and Present), Nobel Tıp Kitabevleri Ltd., Istanbul, 334-335, 1999.
- Sezik E, Yeşilada E, Honda G, Takaishi Y, Takeda Y, Tanaka T. Traditional medicine in Turkey X. Folk medicine in Central Anatolia, *J. Ethnopharmacol.*, 75, 95-115, 2001.
- Anıl H. Elucidation of the structure of saponins from the leaves of *Verbascum nigrum* L., *Doğa Seri A*, 4(1), 1-5, 1980.
- Tschesche R, Sepulveda S, Braun TM. Triterpenes XXXIII. Saponin of *Verbascum phlomoides* blossoms, *Chem. Ber.*, 113(5), 1754-1760, 1980.
- Seifert K, Preiss A, John S, Schmidt J, Lien NT, Lavaud C, Massiot G. Triterpene saponins from *Verbascum songaricum*, *Phytochemistry*, 30(10), 3395-3400, 1991.
- Klimek B, Lavaud C, Massiot G. Saponins from *Verbascum nigrum*, *Phytochemistry*, 31(12), 4368-4370, 1992.
- Tatlı II, Akdemir ZS, Bedir E, Khan IA. Saponin, iridoid, phenylethanoid and monoterpene glycosides from *Verbascum pterocalycinum* var. *mutense* Hub.-Mor., *Turk. J. Chem.*, 28, 111-122, 2004.
- De Pascual Teresa J, Hernandez JM, Diaz F, Grande M. Chemical compounds from *Verbascum lychnitis* L. Lychnitoid, *An. Quim. Ser. C*, 78(1), 108-111, 1982.
- De Pascual Teresa J, Diaz F, Grande M. Components of *Verbascum thapsus* L. III. Contribution to the study of saponins, *An. Quim. Ser. C*, 76(2), 107-110, 1980.
- De Pascual Teresa J, Diaz F, Grande M. Components of *Verbascum thapsus* L. I. Triterpenes, *An. Quim.*, 74(2), 311-314, 1978a.
- Hartleb I, Seifert K. Triterpenoid saponins from *Verbascum songaricum*, *Phytochemistry*, 38(1), 221-224, 1995.
- Miyase T, Horikoshi C, Yabe S, Miyasaka S, Melek FR, Kusano G. Saikosaponin homologs from *Verbascum* spp. The structures of mulleinsaponins I-VII, *Chem. Pharm. Bull.*, 45(12), 2029-2033, 1997.
- Klimek B. Hydroxycinnamoyl ester glycosides and saponins from flowers of *Verbascum phlomoides*, *Phytochemistry*, 43(6), 1281-1284, 1996a.
- Hartleb I, Seifert K. Songarosaponin D-A, triterpenoid saponin from *Verbascum songaricum*, *Phytochemistry*, 35(4), 1009-1011, 1994.
- Hernandez-Hernandez JM. Determinacion de la estructura de los glicosidos de *Verbascum lychnitis* por espectrometria de masas, *Quimica e industria*, 31(7-8), 503-507, 1985.
- Abou Gazar H, Tasdemir D, Ireland CM, Çalis I. Iridoids and triterpene saponins from *Verbascum wiedenmannianum* (Scrophulariaceae), *Biochem. Syst. Ecol.*, 31, 433-436, 2003b.
- Yıldırım Y, Çolak N. A new saponin from *Verbascum ancyritanum* Bornm, *Ankara Ecz. Fak. Der.*, 25(1), 39-42, 1996.
- Eribekyan MI, Arutyunyan LS, Mnatsakanyan VA. Iridoid glycosides of *Verbascum cheiranthifolium*, *Khim. Prir. Soedin.*, (5), 727-728, 1989.
- Agababyan EYu, Arutyunyan LS, Mnatsakanyan VA, Gacs-Baitz E, Radics L. *Verbascum georgicum* iridoids, *Khim. Prir. Soedin.*, (4), 446-451, 1982.
- Akdemir ZS, Tatlı II, Bedir E, Khan IA. Aceylated iridoid glycosides from *Verbascum lasianthum* Boiss., *Turk. J. Chem.*, 28, 101-109, 2004a.
- Agababyan EYu, Arutyunyan LS, Mnatsakanyan VA. Iridoid glycosides of *Verbascum laxum*, *Khim. Prir. Soedin.*, (1), 90-96, 1987.
- Seifert K, Schmidt J, Lien NT, John S. Iridoids from *Verbascum* species, *Planta Med.*, (5), 409-411, 1985.
- Osvath K, Papay V, Toth L. Constituents of the flowers of *Verbascum phlomoides* L. and their therapeutic application, *Herba Hung.*, 21(2-3), 141-147, 1982.
- Mnatsakanyan VA, Arutyunyan LS, Eribekyan MI. Iridoid glycosides of *Verbascum saccatum*, *Khim. Prir. Soedin.*, (1), 38-41, 1983.
- Bianco A, Guiso M, Iavarone C, Passacantilli P, Trogolo C. 6-O-β-D-Xylopyranosylaucubin from *Verbascum sinuatum*, *Phytochemistry*, 19, 571-573, 1980.
- Kalpoutzakis E, Aligiannis N, Mitakou S, Skaltsounis AL. Verbaspinoside, a new iridoid glycoside from *Verbascum spinosum*, *J. Nat. Prod.*, 62, 342-344, 1999.
- Magiatis P, Mitaku S, Tsitsa E, Skaltsounis AL., Harvala C. Verbascoside derivatives and iridoid glycosides from *Verbascum undulatum*. *Nat. Prod. Lett.*, 12(2), 111-115, 1998.
- Pardo F, Perich F, Torres R, Monache FD. Plant irido-

- id glycosides and phyto-growth-inhibitory activity of *Verbascum virgatum*, *Biochem. Syst. Ecol.*, 32, 367-370, 2004.
31. Bianco A, Guiso M, Passacantilli P, Francesconi A. Iridoid and phenylpropanoid glycosides from new sources, *J. Nat. Prod.*, 47(5), 901-902, 1984.
 32. Seifert K, Schöpp E, John S, Hesse M. Iridoide aus *Verbascum nigrum*, *Helv. Chim. Acta*, 65(5), 1678-1685, 1982.
 33. Bianco A, Guiso M, Iavarone C, Passacantilli P, Trogolo C. 6-O- α -Sinuatosylaucubin from *Verbascum sinuatum*, *Phytochemistry*, 20(3), 465-468, 1981a.
 34. Skaltsounis AL, Tsitsa-Tzardis E, Demetzos C, Harvala C. Unduloside, a new iridoid glycoside from *Verbascum undulatum*, *J. Nat. Prod.*, 59 (7), 673-675, 1996.
 35. Swiatek L, Salama O, Sticher O. 6-O- β -D-Xylopyranosylcatalpol, a new iridoid glycoside from *Verbascum thapsiforme*, *Planta Med.*, 45, 153, 1982.
 36. Khuroo MA., Qureshi MA, Razdan TK, Nichols P. Sterones, iridoids and a sesquiterpene from *Verbascum thapsus*, *Phytochemistry*, 27(11), 3541-3544, 1988.
 37. Bianco A, Guiso M, Iavarone C, Passacantilli P, Trogolo C. Sinuatol (6-O- α -L-rhamnopyranosyl-aucubin) from *Verbascum sinuatum*, *Planta Med*, 41, 75-79, 1981b.
 38. Vesper T, Seifert K. Iridoids from *Verbascum nigrum*, *Liebigs Ann. Chem.*, (7), 751-753, 1994.
 39. Magiatis P, Melliou E, Tsitsa E, Charvala C, Mitaku S. Two new acylated iridoid glycosides from *Verbascum undulatum*. *Z. Naturforsch.*, 55c, 667-669, 2000.
 40. Swiatek L. Iridoid glycosides in the Scrophulariaceae family, *Acta Pol. Pharm.*, 30(2), 203-212, 1973.
 41. Serdyuk LI, Dzhumyrko SF., Kompantsev VA. Iridoids and flavonoids of *Verbascum lychnitis*, *Khim. Prir. Soedin.*, (4), 545-546, 1976.
 42. Swiatek L, Luczak S, Grabias B. Occurrence of iridoids in various organs of *Verbascum phlomoides* L. and *Verbascum thapsiforme* Schrad, *Farm. Pol.*, 40(7), 415-418, 1984b.
 43. Tatlı II, Akdemir ZS, Bedir E, Khan IA. 6-O- α -L-Rhamnopyranosylcatalpol derivative iridoids from *Verbascum cilicicum*, *Turk. J. Chem.*, 27, 765-772, 2003.
 44. Eribekyan MI, Arutyunyan LS, Mnatsakanyan VA. Iridoid glycosides of *Verbascum sinuatum*, *Khim. Prir. Soedin.*, (1), 146-147, 1987.
 45. Warashina T, Miyase T, Ueno A. Iridoid glycosides from *Verbascum thapsus* L., *Chem. Pharm. Bull.*, 39(12), 3261-3264, 1991.
 46. Seifert K, Lien NT, Schmidt J, John S, Popov SS, Porzel A. Iridoids from *Verbascum pulverulentum*, *Planta Med.*, 55(5), 470-473, 1989.
 47. Falsone G, Laryea MD, Crea AEG, Finner E. Iridoids from *Verbascum sinuatum*, *Journal of Medicinal Plant Research*, 44, 150-153, 1982.
 48. Klimek B. 6-O-*p*-coumaroylcatalpol from *Verbascum lychnitis*, *Planta Med.*, 57(3), 298, 1991a.
 49. Akdemir ZS, Tatlı II, Bedir E, Khan IA. Iridoid and phenylethanoid glycosides from *Verbascum lasianthum* Boiss., *Turk. J. Chem.*, 28, 227-234, 2004b.
 50. Magiatis P, Spanakis D, Mitaku S, Tsitsa E, Mentis A, Harvala C. Verbalactone, a new macrocyclic dimer lactone from the roots of *Verbascum undulatum* with antibacterial activity, *J. Nat. Prod.*, 64, 1093-1094, 2001.
 51. Arutyunyan LS, Agababyan EYu, Mnatsakanyan VA. Verbascoside B—a minor iridoid from *Verbascum georgicum*, *Khim. Prir. Soedin.*, (5), 573-576, 1983.
 52. Grabias B, Junior P, Swiatek L. Isolation and identification of iridoid glucoside from *Verbascum olympicum* Boiss., *Herba Pol.*, 35(1), 3-8, 1989.
 53. Agababyan EYu, Arutyunyan LS, Mnatsakanyan VA, Gacs-Baitz E, Radics L. Verbascoside from *Verbascum georgicum* Benth., *Arm. Khim. Zh.*, 38(11), 709-712, 1985.
 54. Klimek B. Verbascoside in the flowers of some *Verbascum* species, *Acta Pol. Pharm.*, 48(3-4), 51-54, 1991b.
 55. Tatlı II. Bazı *Verbascum* Türleri Üzerinde Farmakognozik Araştırmalar, Hacettepe Üniversitesi, Sağlık Bilimleri Enstitüsü, Farmakognozik Anabilim Dalı, Doktora Tezi, Ankara, 2004.
 56. Elgindi MR, Mabry TJ, Melek FR, Miyase T. Isoflavones and a phenylethanoid from *Verbascum sinaiticum*, *Asian Journal of Chemistry*, 11(4), 1534-1536, 1999.
 57. Andary C, Wylde R, Laffite C, Privat G, Winternitz F. Structures of verbascoside and orobanchoside, caffeic acid sugar esters from *Orobancha rapum-genitae*, *Phytochemistry*, 21(5), 1123-1127, 1982.
 58. Abou Gazar H, Bedir E, Khan IA, Çalıř I. Wiedemanniosides A-E: new phenylethanoid glycosides from the roots of *Verbascum wiedemannianum*, *Planta Med.*, 69, 814-819, 2003a.
 59. Elgindi MR, Mabry TJ. Phenylethanoid glycosides from *Verbascum sinaiticum*, *Asian Journal of Chemistry*, 12(1), 127-130, 2000.

60. Roussel JL. Thèse Doct. Ès Sci. Pharm., Montpellier, France, 1983.
61. Klimek, B. 6'-O-Apiosyl-verbascoside in the flowers of mullein (*Verbascum speices*), *Acta Pol. Pharm.*, 53(2), 137-140, 1996b.
62. Warashina T, Miyase T, Ueno A. Phenylethanoid and lignan glycosides from *Verbascum thapsus*, *Phytochemistry*, 31(3), 961-965, 1992.
63. Manav Yalcın S. Flavonoids from *Verbascum cheiranthifolium* Boiss., *Marmara Univ. Eczacılık Derg.*, 5(2), 93-97, 1989.
64. Kawashty SA. Flavanoids of *Verbascum* species in Egypt, *Bull. N R C*, 22(1), 11-17, 1997.
65. Papay V, Toth L, Osvath K, Bujtas Gy. Flavonoids from *Verbascum phlomoides* L., *Pharmazie*, 35(5-6), 334-335, 1980.
66. Yuldashev MP. Flavonoids of roots of *Verbascum songaricum*, *Khim. Prir. Soedin.*, 6, 951-952, 1996.
67. Klimek B, Krolikowska M. Flavonoids in flowers of *Verbascum thapsiforme* and *Verbascum phlomoides*, *Acta Pol. Pharm.*, 41(2), 259-264, 1984.
68. Klimek B. Flavonoid glucuronides from *Verbascum lychnitis* and *V. nigrum*, *Acta Pol. Pharm.*, 52(1), 53-56, 1995.
69. Souleles C, Geronikaki A. Flavonoids from *Verbascum thapsus*, *Scientia Pharmaceutica*, 57, 59-61, 1989.
70. Afifi MSA, Ahmed MM, Pezzuto JM, Kinghorn AD. Cytotoxic flavonolignans and flavones from *Verbascum sinaiticum* leaves, *Phytochemistry*, 34(3), 839-841, 1993.
71. Mehrotra R, Ahmed B, Vishwakarma RA, Thakur RS. Verbacoside: a new luteolin glycoside from *Verbascum thapsus*, *J. Nat. Prod.*, 52(3), 640-643, 1989.
72. Naumov P, Kuzmanovski I, Stefova M. Flavonoids of *Verbascum scardicolum* and *Melampyrum scardicum*, *Glas. Hem. Tehnol. Maked.*, 17(1), 41-44, 1998.
73. Tschesche R, Delhvi S, Sepulveda S. Tamarixetin glycosides from the flowers of *Verbascum phlomoides*, *Phytochemistry*, 18(7), 1248-1249, 1979.
74. Kawamo H, Hagiwara Y, Mizuno O, Kumakura M. Isolation of an antihistamine flavone from a plant, *Verbascum thapsus*, *Jpn. Kokai Tokkyo Koho JP 63,227,584 [88,227,584]* (Cl. C07D311/28), 21 Sep 1988, *Appl.* 87/61,304, 18 Mar 1987; 3 pp, 1988.
75. Ulubelen A, Cetin ET, Gueran A. Steroidal compounds and hydrocarbons of *Verbascum lasianthum*, *Planta Med.*, 27(1), 14-17, 1975.
76. Erlaçın S, Gözler B. Sterols and sterol glycosides of *Verbascum pycnostachyum* Boiss. et Heldr., *Pharmazie*, 37, 149-150, 1982.
77. Swiatek L, Dombrowicz E, Kurowska A. Investigation of seed oils from *Verbascum phlomoides* and *V. thapsiforme*, *Herba Pol.*, 31(1-2), 29-33, 1985.
78. Sepulveda S, Delhvi S, Tschesche R. Partial determination of the structure by spectroscopy methods of a new phytosterol isolated from *Verbascum phlomoides* L. flowers, *Contrib. Cient. Tecnol.*, 10(1), 15-21, 1980.
79. Drandarov K. Verbacine and verballocine, novel macrocyclic spermine alkaloids from *Verbascum pseudonobile* Stoj. Et Stef. (*Scrophulariaceae*), *Tetrahedron Lett.*, 36 (4), 617-620, 1995.
80. Drandarov K, Guggisberg A, Hesse M. Macrocyclic spermine alkaloids from *Verbascum*: the (E/Z)-isomeric pairs (-)(S)-verbasitrine/(-)(S)-isoverbasitrine and (+)(S)-verbametrine/(+)(S)-isoverbametrine: isolation, structure elucidation, and synthesis, *Helv. Chim. Acta*, 82, 229-237, 1999.
81. Youhnovski N, Drandarov K, Guggisberg A, Hesse M. Macrocyclic spermine alkaloids from *Verbascum*: isolation, structure elucidation, and synthesis of the (E/Z)-isomeric pairs (S)-verbaskrine/(S)-isoverbasikrine and (S)-verbamekrine/(S)-isoverbamekrine, *Helv. Chim. Acta*, 82, 1185-1194, 1999.
82. Guggisberg A, Drandarov K, Hesse M. Protoverbine, the parent member of a class of macrocyclic spermine alkaloids, *Helv. Chim. Acta*, 83(11), 3035-3042, 2000.
83. Drandarov K, Hesse M. C₁-Derivatives of macrocyclic spermine alkaloids. Verbamedines versus incasines, *Tetrahedron Lett.*, 43(29), 5025-5027, 2002.
84. Ninova P, Abdusamatov A, Yunusov SYu. Alkaloids from *Verbascum nobile*, *Khim. Prir. Soedin.*, 7(4), 540, 1971.
85. Drandarov K. Verballoscenine, the Z isomer of verbascenine from *Verbascum phoeniceum*, *Phytochemistry*, 44(5), 971-973, 1997.
86. Ziyaev R, Abdusamatov A, Yunusov SYu. Alkaloids from *Verbascum songaricum*, *Khim. Prir. Soedin.*, 7(6), 853-854, 1971.
87. Swiatek L, Kurowska A, Rotkiewicz D. Analysis of fatty and phenolic acids in *Verbascum* flowers, *Herba Pol.*, 30(3-4), 173-8, 1984a.
88. De Pascual Teresa J, Diaz F, Grande M. Components of *Verbascum thapsus* L. II. seed oil, *An. Quim.*, 74(12), 1566-1567, 1978b.
89. Kraus J, Franz G. Mucilage polysaccharide from mullein flowers, *Dtsch. Apoth. Ztg.*, 127(13), 665-669, 1987.